arXiv:1205.3105v2 [math.AC] 3 Jan 2013

ON THE CRITICAL IDEALS OF GRAPHS

HUGO CORRALES AND CARLOS E. VALENCIA

Abstract. The critical group of a connected graph G is a finite abelian group isomorphic to the sandpile group of G. In this paper we introduce some determinantal ideals associated with the generalized Laplacian matrix of G, that we call critical ideals. Critical ideals of a graph generalize the critical group in the sense that we can recover the critical group of a graph from their critical ideals. The main results of this article concern the determination of some minimal generator sets and the reduced Gröbner basis for the critical ideals of the complete graphs, the cycles and the paths. Also, we establish a bound between the number of critical ideals that are trivial and the stability and clique numbers of the graph.

1. Introduction

The critical group of a graph (or in general of a digraph) is a finite abelian group that arises in several contexts. Its origin comes from statistical physics under the name of the *abelian sandpile model*, see for instance [6] and [14]. More precisely, the critical group is isomorphic to the sandpile group introduced by Dhar in [14], which generalizes the case of a grid from [6]. In the context of algebraic geometry the critical group is known either as the Picard or the Jacobian group. Also, in the context of combinatorics it is known as the sandpile group or the chip firing game. The critical group has been studied by several authors, for instance see [7, 9, 10, 18, 19, 22, 24]. However, the structure of the critical group has been only completely determined for a few families of graphs.

The critical group of a graph is closely connected with the Laplacian matrix of the graph, as we now explain. Let $G = (V, E)$ be a digraph with n vertices, $m_{(u,v)}$ the number of directed arcs from u to v, and $A(G)$ is the *adjacency matrix* of G given by $A(G)_{u,v} = m_{(u,v)}$. The *Laplacian matrix* of G is given by

$$L(G) = D^+(G) - A(G),$$

where $D^+(G)$ is the diagonal matrix with the out-degrees of the vertices of G in the diagonal entries. Note that the Laplacian matrix is insensitive to loops in the digraph. When G is a simple connected graph their *critical group*, denoted by $K(G)$, is defined as the torsion subgroup of the cokernel of $L(G)$; for instance see [21]. That is,

$$K(G) \oplus \mathbb{Z} = \mathbb{Z}^V / \operatorname{Im} L(G)^t.$$

Originally in the literature the critical group was only defined for graphs, however it can be defined for any matrix with entries in a principal ideal domain; see [13]. Wagner in [25] defined the critical group for a digraph. Although the critical group and the sandpile group are equivalent for graphs, to define the sandpile group for a digraph is needed that the digraph satisfies some technical conditions, see [23]. The critical group has been defined for simplicial complexes, see [16].

2010 *Mathematics Subject Classification.* Primary 13F20; Secondary 13P10, 05C50, 05E99.

Key words and phrases. Critical ideals, Critical group, Digraphs, Graphs, Generalized Laplacian matrix, Determinantal ideal, Gröbner basis, Stability number, Clique number, Cycle, Complete graph.

The first and second authors was partially supported by CONACyT grant 166059, the second author was partially supported by SNI.

The standard method for calculating the critical group of a graph from their Laplacian matrix $L(G)$ consists of reducing $L(G)$ to its Smith Normal Form. Since this reduction depends strongly on the arithmetic relations between the entries of the matrix, finding this reduction is in general very difficult. In order to avoid some of these complications, we propose to study the critical ideals instead of the critical group.

If $X_G = \{x_u \,|\, u \in V(G)\}$ is the set of variables indexed by the vertices of the digraph G, then the *generalized Laplacian matrix* of G, denoted by $L(G, X)$, is given by

$$L(G,X)_{u,v} = \begin{cases} x_u & \text{if } u = v, \\ -m_{(u,v)} & \text{otherwise.} \end{cases}$$

Godsil in [17, 13.9] gives a similar way to define the generalized Laplacian matrix. Furthermore, if $\mathcal{P}$ is a Principal Ideal Domain (PID) with $\mathbb{Z} \subseteq \mathcal{P}$ and $\mathcal{P}[X_G]$ is the polynomial ring over $\mathcal{P}$ in the n variables X_G, then the *critical ideals* of G are the determinantal ideals given by

$$I_i(G,X) = \langle \text{minors}_i(L(G,X)) \rangle \subseteq \mathcal{P}[X_G] \text{ for all } 1 \leq i \leq n,$$

where $\text{minors}_i(L(G,X))$ is the set of determinant of all the i-square submatrices of $L(G,X)$. Note that we can define (without any technical issue) the critical ideals of a $n \times n$ matrix M with entries in $\mathcal{P}$ as $I_i(M,X) = \langle \text{minors}_i(L(M,X)) \rangle \subseteq \mathcal{P}[X]$ for all $i = 1, \ldots, n$, where

$$L(M,X)_{u,v} = \begin{cases} x_u & \text{if } u = v, \\ -M_{u,v} & \text{otherwise.} \end{cases}$$

In [11] is defined a binomial ideal, called the toppling ideal, associated to the Laplacian matrix of a graph, this ideal encode topplings in the abelian sandpile model.

This definition for the critical ideals of a digraph was motivated from the fact that the critical group of G can be calculated in the following way: If G be a graph with n vertices and $K(G) \cong \bigoplus_{i=1}^{n-1} \mathbb{Z}_{f_i}$ with $f_1 | f_2 | \cdots | f_{n-1}$, then

$$\langle \prod_{j=1}^{i} f_j \rangle = \langle \text{minors}_i(L(G)) \rangle.$$

As you can imagine, the critical ideals and the critical group of a graph are closely related, for instance see theorems 3.6 and 3.7. Moreover, critical ideals are very useful to get a better understanding of its critical group. In section 3, we will show that the critical ideals are better behaved than the critical group. For instance, if $\gamma_{\mathcal{P}}(G)$ is the number of critical ideals over the base ring $\mathcal{P}$ that are trivial, then theorem 3.11 claims that

$$\gamma_{\mathcal{P}}(G) \leq 2(|V(G)| - \alpha(G)) \text{ and } \gamma_{\mathcal{P}}(G) \leq 2(|V(G)| - \omega(G)) + 1,$$

where $\alpha(G)$ is the stability number and $\omega(G)$ is the clique number of the graph, respectively. That is, the invariant $\gamma_{\mathcal{P}}$ is closely related to the combinatorics of the graph. Also, if H is an induced subdigraph of G, then $\gamma_{\mathcal{P}}(H) \leq \gamma_{\mathcal{P}}(G)$ at contrast to the behavior of the number of invariant factors of the critical group of induced subdigraphs, see 3.1.

The main goal of this article is the study of the critical ideals of graphs. The main results of this article are contained in sections 3 and 4. Section 3 contains the basic properties of the critical ideals and we define the invariant $\gamma_{\mathcal{P}}$ as the number of the critical ideals that are trivial. Also, in this section we get the reduced Gröbner basis of the complete graph. Finally, we explore the relationship of the critical ideals of a graph with the characteristic polynomial of their adjacency matrix.

The reduced Gröbner basis of the critical ideals of the cycles and some combinatorial expression for the minors of the generalized Laplacian matrix of a digraph are presented in section 4.

2. Preliminaries

A *graph* G is a pair (V, E), with V a finite set and E a collection of unordered pairs of elements of V. The elements of V are called *vertices* and the elements of E are called *edges*. For simplicity, sometimes an edge $e = \{x, y\}$ will be denoted by xy. The sets of two or more edges with the same ends are called *multiple edges*. A *loop* is an edge incident to a unique vertex. A *multigraph* is a graph with multiple edges and without loops.

A *digraph* D is a pair (V, E), where V is a finite set and E is a set of ordered pairs of elements of V. The elements of V and E are called *vertices* and *arcs*, respectively. Given an arc $e = (x, y)$, we say that x is the initial vertex of e and y is the terminal vertex of e. Sometimes for simplicity an arc (u, v) will be denoted by $\overrightarrow{uv}$. The number of arcs with initial vertex x and terminal vertex y will be denoted by $m_{(x,y)}$. The *out-degree* of a vertex x of a digraph D, denoted by $d^+_D(x)$, is the number of arcs in D with initial vertex x. Given a subset U of the vertices of a digraph G, the induced subdigraph by U, denoted by $G[U]$, is the subdigraph of G that has U as vertex set and $E = \{(u, v) \,|\, u, v \in U \text{ and } (u, v) \in E(G)\}$ as arc set. We say that a subdigraph H of G is induced if $H = G[U]$ for some $U \subset V(G)$. In a similar way we can define an induced subgraph.

Note that there exists a natural inclusion of the graphs in the digraphs. Actually, any graph G can be identified with a digraph if we replace each edge $\{u, v\}$ of G by the pair of arcs (u, v) and (v, u). You can consult [15] and [5] for any unexplained concept of graph and digraph theory, respectively.

Now, we will fix some notation for the minors of a matrix. Let $M \in M_n(\mathcal{P})$ be a $n \times n$ matrix with entries on $\mathcal{P}$, $I = \{i_1, \dots, i_r\} \subseteq [n]$, and $J = \{j_1, \dots, j_s\} \subseteq [n]$. The submatrix of M formed by rows $i_1, \dots, i_r$ and columns $j_1, \dots, j_s$ is denoted by $M[I; J]$. On the other hand, the submatrix obtained from M by deleting rows $i_1, \dots, i_r$ and columns $j_1, \dots, j_s$ will be denoted by $M(I; J)$; that is, $M(I; J) = M[I^c; J^c]$. If $|I| = |J| = r$, then $M[I; J]$ is called a *r-square submatrix* or a *square submatrix* of size r of M. Moreover, if $I = J$, then $M[I; J]$ is a *principal square submatrix*. A *r-minor* is the determinant of a r-square submatrix. The set of i-minors of a matrix M will be denoted by $\text{minors}_i(M)$. We say that $M, N \in M_{n\times n}(\mathcal{P})$ are *equivalent*, denoted by $N \sim M$, if there exist $P, Q \in GL_n(\mathcal{P})$ (invertible matrices) such that $N = PMQ$. Is not difficult to see that if $N \sim M$, then $K(M) = \mathcal{P}^n/M^t\mathcal{P}^n \cong \mathcal{P}^n/N^t\mathcal{P}^n = K(N)$.

To finish this section we will recall some useful results on Gröbner basis.

2.1. Gröbner Basis.

Usually the theory of Gröbner basis deals with ideals in a polynomial ring over a field. However, in this paper we will deal with ideals in a polynomial ring over the integers. There exists a theory of Gröbner basis over almost all kind of rings.

We recall some basic concepts on Gröbner basis, see [1]. At first, an *order term* in $\mathcal{P}[x_1, \dots, x_n]$ is a total order $\prec$ in the set of monomials, such that

- **(i):** $1 \prec x^\alpha$ for all $\mathbf{0} \neq \alpha \in \mathbb{N}^n$ and
- **(ii):** If $x^\alpha \prec x^\beta$, then $x^{\alpha+\gamma} \prec x^{\beta+\gamma}$ for all $\gamma \in \mathbb{N}^n$,

where $\mathbf{x}^\alpha = x_1^{\alpha_1} \cdots x_n^{\alpha_n}$.

Now, given an order term $\prec$ and $p \in \mathcal{P}[X]$, let $\mathrm{lt}(p)$ and $\mathrm{lc}(p)$ be the *leading term* and the *leading coefficient* of p, respectively. Given a subset S of $\mathcal{P}[X]$ their leading term ideal of S is the ideal

$$\mathrm{Lt}(S) = \langle \mathrm{lt}(s) \,|\, s \in S \rangle.$$

A set of nonzero polynomials B of an ideal I is called a *Gröbner basis* of I if $\mathrm{Lt}(B) = \mathrm{Lt}(I)$. A good characterization of Gröbner basis is given in terms of S-polynomials.

Definition 2.1. *Let f, f' be polynomials in $\mathcal{P}[X]$ and B be a set of polynomials in $\mathcal{P}[X]$. We say that f reduces strongly to f' modulo B if*

- *$lt(f') \prec lt(f)$ and*
- *there exist $b \in B$ and $h \in \mathcal{P}[X]$ such that $f' = f - hb$.*

Moreover, if $f^* \in \mathcal{P}[X]$ can be obtained from f in a finite number of reductions, we write $f \to_B f^*$.

Given f and g polynomials in $\mathcal{P}[X]$, their *S-polynomial*, denoted by $S(f,g)$, is given by

$$S(f,g) = \frac{c}{c_f}\frac{X}{X_f}\, f - \frac{c}{c_g}\frac{X}{X_g}\, g$$

where $X_f = lt(f)$, $c_f = lc(f)$, $X_g = lt(g)$, $c_g = lc(g)$, $X = \mathrm{lcm}(X_f, X_g)$, and $c = \mathrm{lcm}(c_f, c_g)$.

The next lemma gives us an useful criterion for checking if a set of generators of an ideal is a Gröebner basis.

Lemma 2.2. *Let I be an ideal of polynomials over a PID and B a generating set of I. Then B is a Gröbner basis for I if and only if $S(f,g) \to_B 0$ for all $f \neq g \in B$.*

3. THE CRITICAL IDEALS OF GRAPHS

In this section we will introduce the main concept of this article: the critical ideals of a digraph. We will begin this section by defining of the critical ideals of a digraph, we will present some examples and we will discuss some of their basic properties. Afterwards, we will introduce the number of critical ideals that are trivial as an invariant of the graph and will establish a bound between this invariant and the stability and clique numbers. Also, we will present a minimal set of generators and a reduced Gröbner basis for the critical ideals of the complete graphs. As byproduct we will get expressions for the critical groups for some graphs. Finally, we will explore the relation between the critical ideals of a graph and the characteristic polynomial of its adjacency matrix.

The critical ideals are some determinantal ideals in a polynomial ring, which generalize the critical group and the characteristic polynomial of a graph.

Definition 3.1. *Given a digraph G with n vertices and $1 \leq i \leq n$, let*

$$I_i(G,X) = \langle \mathrm{minors}_i(L(G,X)) \rangle \subseteq \mathcal{P}[X_G]$$

be the i-th critical ideal of G.

Note that in general the critical ideals depend on the base ring $\mathcal{P}$; in this article we are mainly interested when $\mathcal{P} = \mathbb{Z}$. By convention, $I_i(G,X) = \langle 1 \rangle$ if $i \leq 0$ and $I_i(G,X) = \langle 0 \rangle$ if $i > n$. Clearly $I_n(G,X)$ is a principal ideal generated by the determinant of the generalized Laplacian matrix.

Now, we will present an example that illustrate the concept of critical ideals of a digraph.

Example 3.2. *Let G be the complete graph with six vertices minus the perfect matching formed by the edges $M_3 = \{v_1v_4, v_2v_5, v_3v_6\}$ (see figure 1.a) and $\mathcal{P} = \mathbb{Z}$. Then,*

$$L(G,X) = \begin{bmatrix} x_1 & -1 & -1 & 0 & -1 & -1 \\ -1 & x_2 & -1 & -1 & 0 & -1 \\ -1 & -1 & x_3 & -1 & -1 & 0 \\ 0 & -1 & -1 & x_4 & -1 & -1 \\ -1 & 0 & -1 & -1 & x_5 & -1 \\ -1 & -1 & 0 & -1 & -1 & x_6 \end{bmatrix}$$

Using Mathematica, it is not difficult to see that $I_i(G) = \langle 1 \rangle$ for $i = 1, 2$ and

$$I_i(G,X) = \begin{cases} \big\langle 2, x_1, x_2, x_3, x_4, x_5, x_6 \big\rangle & \text{if } i = 3, \\ \big\langle \{x_rx_s \,|\, v_rv_s \in E(G)\} \cup \{2x_r + 2x_s + x_rx_s \,|\, v_rv_s \notin E(G)\} \,\big\rangle & \text{if } i = 4, \\ \big\langle \{x_kx_l(x_r + x_s + x_rx_s) \,|\, (r,s,k,l) \in S(G)\} \cup \{p_{(r,s,k,l)} \,|\, v_rv_s, v_kv_l \notin E(G)\} \big\rangle & \text{if } i = 5, \\ \big\langle x_1x_2x_3x_4x_5x_6 - \sum_{(r,s,k,l)\in S(G)} x_rx_sx_kx_l - 2\sum_{(r,s,k)\in T(G)} x_rx_sx_k \big\rangle & \text{if } i = 6, \end{cases}$$

where

$$S(G) = \{(r,s,k,l) \,|\, v_rv_s \notin E(G),\ v_kv_l \in E(G),\ \text{and } \{i,j\} \cap \{k,l\} = \emptyset\},$$

$T(G)$ are the triangles of G, and $p_{(r,s,k,l)} = (x_r + x_s)(x_k + x_l + x_kx_l) + (x_k + x_l)(x_r + x_s + x_rx_s)$.

Note that the expressions of the critical ideals of G depends heavily on their combinatorics.

Now, let us turn our attention to one of the most basic properties of the critical ideals.

Proposition 3.3. *If G is a digraph, then*

$$\langle 0 \rangle \subsetneq I_n(G) \subseteq \cdots \subseteq I_2(G) \subseteq I_1(G) \subseteq \langle 1 \rangle.$$

Moreover, if H is an induced subdigraph of G with m vertices, then

$$I_k(H) \subseteq I_k(G) \text{ for all } 1 \le k \le m.$$

Proof. Let M be a $k{+}1 \times k{+}1$ matrix with entries on $\mathcal{P}[X_G]$. Since $\det(M) = \sum_{i=1}^{k+1} M_{i,1} \det(M(i;1))$, $I_{k+1}(G) \subseteq I_k(G)$ for all $1 \le k \le n-1$. On the other hand, since any submatrix of $L(H,X)$ is also a submatrix of $L(G,X)$, $\text{minors}_k(L(H,X)) \subseteq \text{minors}_k(L(G,X))$ for all $1 \le k \le n$ and therefore $I_k(H) \subseteq I_k(G)$. □



(a) $K_6 \setminus M_3$ (b) G_1 (c) G_2

FIGURE 1. (a) $K_6 \setminus M_3$, (b) G_1, (c) G_2.

If the digraph is not connected, then we can express its critical ideals as a function of the critical ideals of their connected components.

Proposition 3.4. *If G and H are vertex disjoint digraphs, then*

$$I_i(G \sqcup H) = \left\langle \bigcup_{j=0}^{i} I_j(G) I_{i-j}(H) \right\rangle \text{ for all } 1 \leq i \leq |V(G \sqcup H)|.$$

Proof. Let $Q = M \oplus N$ where $M \in M_m(B)$ and $N \in M_n(B)$, $k \in [m+n]$, and $\mathbf{r}, \mathbf{s} \subseteq [m+n]$ with $|\mathbf{r}| = |\mathbf{s}| = k$. It follows by induction on $|\mathbf{r}_m|$, that

$$\det(Q[\mathbf{r};\mathbf{s}]) = \begin{cases} \det(Q[\mathbf{r}_m;\mathbf{s}_m]) \cdot \det(Q[\mathbf{r}_n;\mathbf{s}_n]) & \text{if } |\mathbf{r}_m| = |\mathbf{s}_m|, \\ 0 & \text{otherwise,} \end{cases}$$

where $\mathbf{r}_m = \mathbf{r} \cap [m]$, $\mathbf{s}_m = \mathbf{s} \cap [m]$, $\mathbf{r}_n = \mathbf{r} \setminus \mathbf{r}_m$, and $\mathbf{s}_n = \mathbf{s} \setminus \mathbf{s}_m$.

Now, since $L(G \sqcup H, X) = L(G, X) \oplus L(H, X)$, we get that $\text{minors}_i(L(G \sqcup H, X)) \setminus 0 \subseteq \{m_1 \cdot m_2 \,|\, m_1 \in \text{minors}_j(L(G,X)) \text{ and } m_2 \in \text{minors}_{i-j}(L(H,X)) \text{ for some } 0 \leq j \leq i\}$ for all $1 \leq i \leq |V(G \cup H)|$ and the result is obtained. ☐

Let T_v be the trivial graph composed by the vertex v. Since $I_1(T_v) = \langle x_v \rangle$, applying proposition 3.4 we get the critical ideals of the trivial graph with n vertices.

Corollary 3.5. *If $n \geq 1$ and T_n is the graph with n isolated vertices, then*

$$I_i(T_n) = \big\langle \{\prod_{j \in J} x_j \,\big|\, |J| = i\} \big\rangle \text{ for all } 1 \leq i \leq n.$$

Note that the critical ideals of T_n do not depend on the base ring.

Now, we will establish some basic relationships between the critical ideals and the critical group. Before to doing this we will introduce some notation. Given a digraph G with n vertices and $\mathbf{d} \in \mathcal{P}^{V(G)}$, let $L(G, \mathbf{d})$ be the matrix obtained from $L(G, X)$ where we put $x_v = \mathbf{d}_v$ for all $v \in V(G)$. Also, let

$$I_i(G, \mathbf{d}) = \{f(\mathbf{d}) \,|\, f \in I_i(G, X)\} \text{ for all } 1 \leq i \leq n$$

be ideals in $\mathbb{Z}$. Given an induced subdigraph H of G, the *degree vector* of H in G is given by $d_G(H)_v = \deg^+_G(v)$ for all $v \in V(H)$.

Theorem 3.6. *If $\mathcal{P} = \mathbb{Z}$, G is a digraph with n vertices, and $K(G) \cong \bigoplus_{i=1}^{n-1} \mathbb{Z}_{f_i}$ with $f_1 | \cdots | f_{n-1}$, then*

$$I_i(G, d_G(G)) = \langle \prod_{j=1}^{i} f_j \rangle \text{ for all } 1 \leq i \leq n-1.$$

Proof. Clearly $L(G, d_G(G)) = L(G)$, the usual Laplacian matrix of G. Thus $\text{minors}_i(L(G, d_G(G))) = \text{minors}_i(L(G))$ for all $1 \leq i \leq n$ and therefore

$$I_i(G, d_G(G)) = \langle \text{minors}_i(L(G, d_G(G))) \rangle = \langle \text{minors}_i(L(G)) \rangle = \langle \prod_{j=1}^{i} f_j \rangle \text{ for all } 1 \leq i \leq n-1.$$

☐

On the other hand, if v is a vertex of G and $L(G, v)$ is the *reduced Laplacian matrix*, given by the matrix obtained from $L(G)$ by removing the row and column v, then we have the following strong version of theorem 3.6:

Theorem 3.7. *Let $\mathcal{P}=\mathbb{Z}$, G be a connected graph (possible with multiple edges) with n vertices, and v a vertex of G. If $K(G)\cong\bigoplus_{i=1}^{n-1}\mathbb{Z}_{f_i}$ with $f_1|\cdots|f_{n-1}$, then*

$$I_i(G\setminus v, d_G(G\setminus v))=\langle\prod_{j=1}^{i} f_j\rangle \text{ for all } 1\leq i\leq n-1.$$

Proof. Since $L(G\backslash v, d_G(G\backslash v))=L(G,v)$ (the usual reduced Laplacian matrix of G), by theorem 3.6 we only need to prove that $K(G)\cong\mathbb{Z}^{V(G)\backslash v}/\mathrm{Im}\, L(G,v)^t$. Now, if $I_{n-1,n-1}$ is the identity matrix of size $n-1$, then

$$\begin{bmatrix} 1 & \mathbf{1} \\ \mathbf{0} & I_{n-1,n-1}\end{bmatrix} L(G) \begin{bmatrix} 1 & \mathbf{1} \\ \mathbf{0} & I_{n-1,n-1}\end{bmatrix}^t = 0\oplus L(G,v).$$

Since $\det(\begin{bmatrix} 1 & \mathbf{1} \\ \mathbf{0} & I_{n-1,n-1}\end{bmatrix})=1$, then $L(G)\sim 0\oplus L(G,v)$ and we get the result. □

Remark 3.8. *Note that, in general theorem 3.7 is not valid for digraphs. However, we can get a similar result for matrices in $M_{n\times n}$ that are symmetric with rank $n-1$ and such that $M\mathbf{1}=\mathbf{0}$.*

The next example shows how theorem 3.7 can be used to recover the critical group of a graph from their critical ideals. In this sense, critical ideals generalize the critical group of a graph.

Example 3.9. *Let H be the complete graph with six vertices minus a perfect matching as in figure 1.a. and G be a graph such that $H=G\setminus v$ for some vertex v of G. Thus, applying theorem 3.7 we can get the critical group of G as an evaluation of the critical ideals of H. For instance, if G_1 is the graph obtained from H (see figure 1.b) by adding a new vertex v and the edges vv_1, vv_3, vv_4, vv_6, then $d_G(H)=(5,4,5,5,4,5)$. Moreover, using the critical ideals of H calculated in the example 3.2, we get that $f_i=1$ for all $i\leq 4$, $f_5=20$, $f_6=140$; that is,*

$$K(G_1)\cong\mathbb{Z}_{20}\oplus\mathbb{Z}_{140}.$$

On the other hand, if we only know the critical ideals of induced subgraphs of G that are different to $G\backslash v$, then we can not reconstruct completely the critical group of G. For instance, if G_2 is the graph obtained by adding the vertices v_5, v_6 and the edges $v_5v_1, v_5v_2, v_6v_2, v_6v_3, v_6v_4$ to the complete graph with four vertices (see figure 1.c), then it is not difficult to see using Mathematica that $f_i(G_2)=1$ for $1\leq i\leq 4$ and $f_5(G_2)=185$. However, when we apply theorem 3.7 to the critical ideals of the induced subgraph by the vertices v_1, v_2, v_3, and v_4 of G_2 (isomorphic to K_4) we can only obtain that $f_1(G_2)=f_2(G_2)=1$, $f_3(G_2)\,|\,5$ and $f_4(G_2)\,|\,175$.

3.1. **The invariant γ.**

In this subsection we will present an invariant that will play an important role on the study of the critical ideals and the critical group of a digraph.

Definition 3.10. *Given a digraph G and a PID $\mathcal{P}$, let $\gamma_{\mathcal{P}}(G)=\max\{i\,|\,I_i(G,X)=\langle 1\rangle\}$.*

There exists a close relation between $\gamma_{\mathcal{P}}(G)$ and the number of invariant factor of the critical group of G that are equal to 1. For instance, if $IF_1(G)$ denote the number of invariant factor of the critical group of G that are equal to 1, then $\gamma_{\mathbb{Z}}(G)\leq IF_1(G)$. We found that $\gamma_{\mathcal{P}}(G)$ behaves better than the number of invariant factors of the critical group of a digraph that are one. For instance, it is not difficult to see from the definition and proposition 3.3 that if H is an induced subdigraph of G, then $\gamma_{\mathcal{P}}(H)\leq\gamma_{\mathcal{P}}(G)$. However, if $n\geq 3$, $G=K_{2,n}$ is a complete bipartite graph, and $H=K_{1,n}$ is an induced subgraph of G, then $IF_1(G)=2<n=IF_1(H)$.

Now we will present a relation between $\gamma_{\mathcal{P}}(G)$ and the stability and the clique numbers of G. Before doing this, we will define the stability and the clique numbers of a graph.

A subset S of the vertices of a graph G is called *stable* or *independent* if does not exist an edge of G with ends in S. A stable set is called *maximal* if it is under the inclusion of sets. The *stability number* of G, denoted by $\alpha(G)$, is given by

$$\alpha(G) = \max\{|S| \,|\, S \text{ is a stable set of } G\}.$$

In a similar way, a subset C of the vertices of a graph G is called *clique* if all the pairs of vertices in C are joined by an edge of G. A clique set is called *maximal* if it is under the inclusion of sets. The *clique number* of G, denoted by $\omega(G)$, is given by

$$\omega(G) = \max\{|C| \,|\, C \text{ is a clique set of } G\}.$$

Theorem 3.11. *If G is a simple multigraph with n vertices, then*

$$\gamma_{\mathcal{P}}(G) \leq 2(n-\omega(G))+1 \text{ and } \gamma_{\mathcal{P}}(G) \leq 2(n-\alpha(G)).$$

Proof. We begin with a simple relation between the critical ideals of G and $G\setminus v$ for some vertex v of G.

Claim 3.12. *If G is a multigraph with $V(G)=\{v_1,\dots,v_n\}$, then*

$$I_j(G,X) \subseteq \langle x_1 I_{j-1}(G\setminus v_1, X\setminus x_1), I_{j-2}(G\setminus v_1, X\setminus x_1)\rangle \text{ for all } 1\leq j\leq n.$$

Proof. Let $I=\{i_1,...,i_j\}\subseteq[n]$, $I'=\{i'_i,...,i'_j\}\subseteq[n]$ and $m_{I,I'}=\det(L(G,X)[I,I'])\in I_j(G)$. If $1\notin I\cup I'$, then $m_{I,I'}\in I_j(G\setminus v_1, X\setminus x_1)$. In a similar way, if $1\in I\Delta I'$, then $m_{I,I'}\in I_{j-1}(G\setminus v_1,X\setminus x_1)$. On the other hand, if $1\in I\cap I'$, then

$$m_{I,I'}\in\langle x_1 I_{j-1}(G\setminus v_1,X\setminus x_1), I_{j-2}(G\setminus v_1,X\setminus x_1)\rangle.$$

Finally, the result follows because $I_j(G\setminus v_1,X\setminus x_1)\subseteq I_{j-1}(G\setminus v_1,X\setminus x_1)\subseteq I_{j-2}(G\setminus v_1,X\setminus x_1)$. □

Now, let $g=\gamma_{\mathcal{P}}(G\setminus v_1)$. Since $I_i(G\setminus v_1,X\setminus x_1)\neq\langle 1\rangle$ for all $i\geq g+1$, by claim 3.12

$$I_{g+3}(G,X)\subseteq\langle x_1 I_{g+2}(G\setminus v_1,X\setminus x_1), I_{g+1}(G\setminus v_1,X\setminus x_1)\rangle\neq\langle 1\rangle$$

and therefore $I_{g+3}(G,X)\neq\langle 1\rangle$. That is, $\gamma_{\mathcal{P}}(G)-\gamma_{\mathcal{P}}(G\setminus v_1)\leq 2$.

Finally, the result follows by using that $\gamma_{\mathcal{P}}(T_{\alpha(G)})=0$ (corollary 3.5), $\gamma_{\mathcal{P}}(K_{\omega(G)})=1$ (theorem 3.14) and the fact that $\gamma_{\mathcal{P}}(G)-\gamma_{\mathcal{P}}(G\setminus v)\leq 2$ for all $v\in V(G)$. □

This result is interesting when either the stability or the clique number are almost the number of vertices of the graph. For instance, if G is the complete graph ($\omega(G)=n$), then $\gamma_{\mathcal{P}}(G)\leq 1$. Theorem 3.14 proves that this bound is tight in this case. Similarly to theorem 3.11, in [21] an upper bound for the number of invariant factor different to one of the critical group of a graph G in terms of the number of independent cycles of G was found.

In [20] a result similar to the one obtained in the proof of theorem 3.11 was obtained. Namely in [20] it was shown that if G is a simple graph and $e\in E(G)$, then the number of invariant factor of G and $G\setminus e$ differ by at most 1.

Clearly, a simple graph G has $\gamma_{\mathcal{P}}(G)=0$ if and only if G is the trivial graph. Moreover, in [2] it was shown that a simple graph G has $\gamma_{\mathbb{Z}}(G)=1$ if and only if G is the complete graph. Also, all the simple graphs with $\gamma_{\mathbb{Z}}$ equal to 2 and 3 were characterized in [2] and [3] .

It is not difficult to prove that the bound $\gamma_{\mathcal{P}}(G)\leq 2(n-\alpha(G))$ is tight; for instance, is easy to prove that if P_{2n+1} is an odd path (see corollary 4.10), then $\alpha(P_{2n+1})=n+1$ and $\gamma_{\mathcal{P}}(P_{2n+1})=2n=2(2n+1-(n+1))$. Moreover, in [12] it was shown that if T is a tree, then γ is equal to the

2-matching number of the tree. This result proves that the bound $\gamma_{\mathcal{P}}(G) \leq 2(n - \alpha(G))$ is tight for any value of the stability number and the number of vertices of the graph. An interesting open question is the characterization of the simple graphs that satisfy the bounds given in theorem 3.11.

Now, we will turn our attention to the critical ideals of the first non-trivial graph.

3.2. **The critical ideals of the complete graphs.** We begin this subsection with an expression for the determinant of the complete graph.

Theorem 3.13. *If K_n is the complete graph with $n \geq 1$ vertices, then*

$$\det(L(K_n, X)) = \prod_{j=1}^{n}(x_j + 1) - \sum_{i=1}^{n}\prod_{j\neq i}(x_j + 1).$$

Proof. We will use induction on n. For $n = 1$, it is clear that $\det(L(K_n, X)) = x_1 = (x_1 + 1) - 1$. Now, assume that $n \geq 2$. Expanding the determinant of $L(K_{n+1}, X)$ by the last column and using induction hypothesis

$$\begin{aligned}
\det(L(K_{n+1}, X)) &= x_{n+1} \cdot \det(L(K_n, X)) + \sum_{k=1}^{n} \det(L(K_n, X))_{x_k=-1} \\
&= x_{n+1} \cdot \left[\prod_{j=1}^{n}(x_j + 1) - \sum_{i=1}^{n}\prod_{j\neq i}(x_j + 1)\right] - \sum_{k=1}^{n}\prod_{j\neq k}(x_j + 1) \\
&= \prod_{j=1}^{n+1}(x_j + 1) - \prod_{j=1}^{n}(x_j + 1) - (x_{n+1} + 1)\sum_{i=1}^{n}\prod_{j\neq i}(x_j + 1) \\
&= \prod_{j=1}^{n+1}(x_j + 1) - \sum_{i=1}^{n+1}\prod_{j\neq i}(x_j + 1).
\end{aligned}$$

□

The next result gives us a description of a reduced Gröbner basis of the critical ideals of the complete graph.

Theorem 3.14. *If K_n is the complete graph with $n \geq 2$ vertices and $1 \leq m \leq n - 1$, then*

$$B_m = \Big\{ \prod_{i \in I}(x_i + 1) \,|\, I \subseteq [n] \text{ and } |I| = m - 1 \Big\}$$

is a reduced Gröbner basis of $I_m(K_n, X)$ with respect to the graded lexicographic order.

Proof. At first, we will prove that B_m generates $I_m(K_n, X)$. If M is a square submatrix of $L(K_n, X)$ of size m, then there exist $I \subset [n]$ with $|I| = m$ and $J \subseteq I$ such that M is equal to

$$L(K_n, X)[I; I]_{\{x_j=-1 \text{ for all } j\in J\}}.$$

Thus, since

$$\det(L(K_n, X))[I; I]_{\{x_j=-1 \text{ for all } j\in J\}} = \begin{cases} \prod_{i_j \in I\setminus J}(x_{i_j} + 1) & \text{if } |J| = 1, \\ 0 & \text{if } |J| \geq 2, \end{cases}$$

and $\det(L(K_n, X))[I; I] = -(x_{i_m}+1)\cdot\det(L(K_n, X))[I; I]_{\{x_{i_m}=-1\}} + \sum_{k=1}^{m} \det(L(K_n, X))[I; I]_{\{x_{i_k}=-1\}}$, then B_m generates $I_m(K_n, X)$.

Finally, we will prove that B_m is a reduced Gröbner basis of $I_m(K_n, X)$ for all $1 \leq m \leq n-1$. Let $I_1, I_2 \subset [n]$ with $|I_1|, |I_2| = m-1$ and $p_I = \prod_{i\in I}(x_i+1)$. It is not difficult to see that $lt(p_I) = \prod_{i\in I} x_i$ and

$$p_{I_1} \cdot \prod_{i\in I_2\setminus I_1} (x_i+1) - p_{I_2} \cdot \prod_{i\in I_1\setminus I_2} (x_i+1) = 0.$$

Thus

$$\begin{aligned} S(p_{I_1}, p_{I_2}) &= \frac{lt(p_{I_2})}{lt(p_{I_1\cap I_2})} p_{I_1} - \frac{lt(p_{I_1})}{lt(p_{I_1\cap I_2})} p_{I_2} \\ &= \left(\prod_{i\in I_2\setminus I_1} (x_i+1) - \prod_{i\in I_2\setminus I_1} x_i \right) \cdot p_{I_2} - \left(\prod_{i\in I_1\setminus I_2} (x_i+1) - \prod_{i\in I_1\setminus I_2} x_i \right) \cdot p_{I_1} \end{aligned}$$

and $S(p_{I_1}, p_{I_2}) \to_{B_m} 0$. Therefore B_m is a reduced Gröbner basis of $I_i(K_n, X)$. □

Remark 3.15. *Note that the critical ideals of K_n do not depend on the base ring. Also, note that, since $\prod_\emptyset = 1$, $I_1(K_n, X) = \langle 1 \rangle$ and therefore $\gamma(K_n) = 1$.*

Using the expression for the critical ideals of the complete graph given in theorem 3.14, we can get the primary decomposition of the critical ideals of the complete graph.

Corollary 3.16. *If K_n is the complete graph with $n \geq 2$ vertices and $1 \leq m \leq n-1$, then*

$$I_m(K_n, X) = \bigcap_{I\subset[n], |I|=n-m+2} \langle \{x_i + 1 \,|\, i \in I\} \rangle.$$

As an application of theorem 3.14 we find the critical group of all the graphs such that $\omega(G) = |V(G)| - 1$.

Corollary 3.17. *Let $n > m \geq 1$ and $K_{n+1} \setminus S_m$ be the graph obtained from K_{n+1} by deleting the m edges of the star S_m. Then*

$$K(K_{n+1} \setminus S_m) = \begin{cases} \mathbb{Z}_{n+1}^{n-2m} \oplus \mathbb{Z}_{n(n+1)}^{m-2} \oplus \mathbb{Z}_{n(n+1)(n-m)} & \text{if } m \leq \lfloor n/2 \rfloor, \\ \\ \mathbb{Z}_{n}^{2m-n} \oplus \mathbb{Z}_{n(n+1)}^{n-m-2} \oplus \mathbb{Z}_{n(n+1)(n-m)} & \text{if } m \geq \lfloor n/2 \rfloor. \end{cases}$$

Proof. If v_{n+1} is the vertex of $K_{n+1} \setminus S_m$ of degree $n-m$, then

$$\mathbf{d} = d_{K_{n+1}-S_m}(K_{n+1} \setminus v_{n+1}) = (\underbrace{n-1, \ldots, n-1}_{m \text{ times}}, \underbrace{n, \ldots, n}_{n-m \text{ times}}).$$

By theorem 3.13, $I_n(K_n, \mathbf{d}) = \langle n^{m-1}(n+1)^{n-m-1}(n-m) \rangle$,

$$I_{i+1}(K_n, \mathbf{d}) = \begin{cases} \langle 1 \rangle & \text{if } i \leq m, \\ \langle (n+1)^{i-m} \rangle & \text{if } m < i \leq n-m, \\ \langle n^{i+m-n}(n+1)^{i-m} \rangle & \text{if } n-m < i < n-1, \end{cases}$$

when $m \leq \lfloor n/2 \rfloor$, and

$$I_{i+1}(K_n, \mathbf{d}) = \begin{cases} \langle 1 \rangle & \text{if } i \leq n-m, \\ \langle n^{i+m-n} \rangle & \text{if } n-m < i \leq m, \\ \langle n^{i+m-n}(n+1)^{i-m} \rangle & \text{if } m < i < n-1, \end{cases}$$

when $m \geq \lfloor n/2 \rfloor$. Finally, applying theorem 3.7 we get the result. □

To our knowledge, the critical group of this graphs had not been calculated before. We finish this section by exploring a relation between the critical ideals and the characteristic polynomial of the adjacency matrix of a graph.

3.3. **The critical ideals and the characteristic polynomials.** In [22], Lorenzini shown a deep relation between the critical group and the Laplacian spectrum of a graph. For instance, Lorenzini (Proposition 3.2) prove that if λ is an integer eigenvalue of $L(G)$ of multiplicity $\mu(\lambda)$, then $K(G)$ contains a subgroup isomorphic to $\mathbb{Z}_\lambda^{\mu(\lambda)-1}$. In this subsection we will present the relationship that exists between the critical ideals and the characteristic polynomial of the adjacency matrix of a graph. If G is loopless and we take $x_i = t$ for all $1 \leq i \leq n$, then $\det(L(G,X))$ is equal to the characteristic polynomial $p_G(t)$ of the adjacency matrix of G and the critical ideals of G are ideals in $\mathcal{P}[t]$. The ideals $I_i(G,t) = I_i(G,X)_{\{x_i=t \,|\, \forall\, 1\leq i\leq n\}}$ are called the *t-critical ideals* of G. It is not difficult to see that if $\mathcal{P}$ is a field, then the critical ideals of G are principal. That is, there exist $p_i(t) \in \mathcal{P}[t]$ for all $1 \leq i \leq n$ such that

$$I_i(G,t) = \langle \prod_{j=1}^{i} p_j(t) \rangle \text{ for all } 1 \leq i \leq n.$$

Thus, $p_G(t) = \prod_{j=1}^{n} p_j(t)$ is a factorization of the characteristic polynomial of the adjacency matrix of G.

Therefore the critical ideals of a graph are a generalization of the characteristic polynomial the adjacency matrix of G. For instance, if G is the complete graph with six vertices minus a perfect matching (as in example 3.2) and $\mathcal{P} = \mathbb{Q}$, then

$$I_i(G,t) = \begin{cases} \langle 1 \rangle & \text{if } 1 \leq i \leq 4, \\ \langle t^2(t+2) \rangle & \text{if } i = 5, \\ \langle t^3(t+2)^2(t-4) \rangle & \text{if } i = 6. \end{cases}$$

Note that in general the t-critical ideals depend on the base ring. For instance, if $\mathcal{P} = \mathbb{Z}$, then

$$I_i(G,t) = \begin{cases} \langle 1 \rangle & \text{if } i = 1,2, \\ \langle 2, t \rangle & \text{if } i = 3, \\ \langle t^2, 4t \rangle & \text{if } i = 4, \\ \langle t^3(t+2), 4t^2(t+2) \rangle & \text{if } i = 5, \\ \langle t^3(t+2)^2(t-4) \rangle & \text{if } i = 6. \end{cases}$$

The critical and t-critical ideals are a stronger invariant than the adjacency spectrum of the graph. For instance, if $\mathcal{P} = \mathbb{Z}$, G_1 and G_2 are the graphs given in figure 2, then they have the same



FIGURE 2. The graphs G_1 and G_2 have the same characteristic polynomial.

adjacency spectrum, but their t-critical ideals are given by

$$I_i(G_1,t)=\begin{cases}\langle 1\rangle & \text{if } 1\leq i\leq 4,\\ \langle 2(t+1),(t+1)\cdot(t^2+1)\rangle & \text{if } i=5,\\ \langle (t-1)\cdot(t+1)^2\cdot(t^3-t^2-5t+1)\rangle & \text{if } i=6,\end{cases}$$

and

$$I_i(G_2,t)=\begin{cases}\langle 1\rangle & \text{if } 1\leq i\leq 3,\\ \langle 2,(t+1)\rangle & \text{if } i=4,\\ \langle 4(t+1),(t+1)\cdot(t-3)\rangle & \text{if } i=5,\\ \langle (t-1)\cdot(t+1)^2\cdot(t^3-t^2-5t+1)\rangle & \text{if } i=6.\end{cases}$$

4. Critical ideals of the cycle

The main goal of this section is to get a minimal set of generators and the reduced Gröbner basis of the critical ideals of the cycle. Before doing this, we will get some combinatorial expressions for the minors of the generalized Laplacian matrix of a digraph. This expression will be very useful in order to get some algebraic relations between the generators of the critical ideals of the cycle.

We begin with some concepts of digraphs. Given a digraph D, a subdigraph C of D is called a *directed* 1*-factor* if and only if $d^+_C(v)=d^-_C(v)=1$ for all $v\in V(C)$. The number of connected components of C will be denoted by $c(C)$.

Theorem 4.1. *If D is a digraph with n vertices (possibly with multiple arcs and loops), then*

$$\det(-A(D))=\sum_{C\in\overrightarrow{\mathcal{F}}}(-1)^{c(C)},$$

where $\overrightarrow{\mathcal{F}}$ is the set of spanning directed 1*-factors of D. Moreover,*

$$\det(L(D,X))=\sum_{U\subseteq V(D)}\det(-A(D[U]^{wl}))\cdot\prod_{v\notin U}x_v,$$

where $D[U]$ is the induced subgraph of D by U and $D[U]^{wl}$ is the graph obtained from $D[U]$ when we delete the possible loops.

Proof. It follows from arguments similar to those in [8][Proposition 7.2]. □

Remark 4.2. *If we identify a graph G with the digraph D_G, obtained from G by replacing each edge uv of G by the arcs $\overrightarrow{uv}$ and $\overrightarrow{vu}$, then an elementary graph is identified with a spanning directed* 1*-factor and therefore the first part of theorem 4.1 is equivalent to* [8]*[Proposition 7.2]. In this way, theorem 4.1 is a generalization of the expression obtained in* [8]*[Proposition 7.2].*

Next we will present an example of the use of theorem 4.1.

Example 4.3. *If D is the digraph given by $V(D)=\{v_1,v_2,v_3,v_4\}$ (see figure 3) and*

$$E(D)=\{\overrightarrow{v_1v_1},\overrightarrow{v_1v_2},\overrightarrow{v_2v_3},\overrightarrow{v_3v_4},\overrightarrow{v_4v_1},\overrightarrow{v_4v_2}\},$$

then the digraph D has two spanning directed 1*-factors; $\{\overrightarrow{v_1v_2},\overrightarrow{v_2v_3},\overrightarrow{v_3v_4},\overrightarrow{v_4v_1}\}$ and $\{\overrightarrow{v_1v_1},\overrightarrow{v_2v_3},\overrightarrow{v_3v_4},\overrightarrow{v_4v_2}\}$. Thus $\det(A(D))=(-1)^1+(-1)^2=0$. Also, since $\{\overrightarrow{v_1v_2},\overrightarrow{v_2v_3},\overrightarrow{v_3v_4},\overrightarrow{v_4v_1}\}$ and $\{\overrightarrow{v_2v_3},\overrightarrow{v_3v_4},\overrightarrow{v_4v_2}\}$ are the directed* 1*-factors of D^{wl},*

$$\det(L(D,X))=x_1x_2x_3x_4-x_1-1.$$



FIGURE 3. A digraph with four vertices.

When G is a tree, the expression of the determinant of the generalized Laplacian given in theorem 4.1 only depends on their matchings. Therefore, in this case we can get an explicit combinatorial expression for the determinant of the generalized Laplacian. Note that, $V(\mu)$ denoted the set of vertex of the induced subgraph by μ.

Theorem 4.4. *If T is a tree, then*

$$\det(L(T,X)) = \sum_{\mu\in\mathcal{V}_1(T)} (-1)^{|\mu|} \prod_{v\notin V(\mu)} x_v.$$

Proof. It follows from theorem 4.1 and the fact that if S is a forest, then

$$\det(A(S)) = \begin{cases} (-1)^{|\mu|} & \text{if } \mu \text{ is a perfect matching of } S, \text{ and} \\ 0 & \text{otherwise.} \end{cases}$$ □

Moreover, theorems 4.1 and 4.4 can be used to get combinatorial expressions for the determinants of the generalized Laplacian matrices of graphs.

Corollary 4.5. *Let n be a natural number, K_n be the complete graph with n vertices, P_n be the path with n vertices, and C_n the cycle with n vertices. Then*

- **(i):** $\det(L(K_n,X)) = \sum_{I\subseteq[n]} (|I|-n+1)\cdot \prod_{i\in I} x_i,$
- **(ii):** $\det(L(P_n,X)) = \sum_{\mu\in\mathcal{V}_1(P_n)} (-1)^{|\mu|}\cdot \prod_{v\notin V(\mu)} x_v,$
- **(ii):** $\det(L(C_n,X)) = \sum_{\mu\in\mathcal{V}_1(C_n)} (-1)^{|\mu|}\cdot \prod_{v\notin V(\mu)} x_v - 2,$

where $\mathcal{V}_1(G)$ is the set of matchings of G.

Proof. (i) It follows by theorem 4.1 and the fact that $\det(A(K_m)) = -m+1$ for all $m \geq 2$. (ii) It follows directly from theorem 4.4. (iii) Let P_i^l be the induced path of C_n of length l that begins in the vertex i of C_n. Since $\mathcal{V}_1(C_n) = \mathcal{V}_1(C_n \setminus v_1v_n) \sqcup \{\mu\in\mathcal{V}_1(C_n)\,|\,v_1v_n\in\mu\}$,

$$\det(L(C_n,X)) = \det(L(P_1^n,X)) - \det(L(P_2^{n-2},X)) - 2 = \sum_{\mu\in\mathcal{V}_1(C_n)} (-1)^{|\mu|}\cdot \prod_{v\notin V(\mu)} x_v - 2.$$ □

Now, we will show that the minors of a generalized Laplacian of a digraph are equal to an evaluation of the determinant of the generalized Laplacian of some digraphs.

Given a digraph (or a graph) D and $u \neq v$ two vertices of D, let $D(u;v)$ be the digraph obtained from D by deleting the arcs leaving u and entering v (remember that each edge of a graph is considered as two arcs in both directions) and identifying the vertices u and v in a new vertex, denoted by $u\circ v$. Also, if $u = v$, then $D(u;v)$ is defined as $D \setminus u$.

Example 4.6. *If P_n is the path with n vertices (simple graph), then*

$$P_n(v_1;v_n)\cong C_{n-1}\setminus\{\overrightarrow{v_1v_2},\overrightarrow{v_{n-1}v_1}\}.$$

Now, given two matrices $M,N\in M_{m\times m}(\mathcal{P})$ we say that M and N are *strong equivalent*, denoted by $N\approx M$, if there exist P and Q permutation matrices such that $N=PMQ$. Also, given $U=\{u_1,\dots,u_s\}$ and $V=\{v_1,\dots,v_s\}$ ordered subsets of $V(D)$ with $s\geq 2$ we define $D(U;V)$ inductively as $D(U\setminus u_s;V\setminus v_s)(u_s;v_s)$.

Lemma 4.7. *If D is a digraph and $U=\{u_1,\dots,u_s\}$, $V=\{v_1,\dots,v_s\}$ ordered subsets of $V(D)$, then*

$$L(D,X)(U;V)\approx L(D(U;V),X)_{\{x_{u_i\circ v_i}=-m_{(v_i,u_i)}\,|\,i=1,\dots,s\}}.$$

Proof. It follows from the construction of $D(U;V)$. □

Remark 4.8. *Clearly, $D(U;V)$ and $L(D(U;V),X)$ depend on the order of the elements of the subsets U and V. However, if $\sigma\in S_s$ is a permutation, then*

$$L(D(U;V),X)_{\{x_{u_i\circ v_i}=-m_{(v_i,u_i)}\,|\,i=1,\dots,s\}}\approx L(D(U;\{v_{\sigma(1)},\dots,v_{\sigma(s)}\}),X)_{\{x_{u_i\circ v_{\sigma(i)}}=-m_{(v_{\sigma(i)},u_i)}\,|\,i=1,\dots,s\}},$$

that is, in some sense $L(D(U;V),X)_{\{x_{u_i\circ v_i}=-m_{(v_i,u_i)}\,|\,i=1,\dots,s\}}$ does not depend on the order of the elements of the subsets U and V.

Example 4.9. *Let D be the cycle with six vertices, $U=\{v_1,v_2\}$, and $V=\{v_6,v_5\}$. Then we get that $D(U;V)$ is the digraph with four vertices $\{v_3,v_4,v_2\circ v_5,v_1\circ v_6\}$ (see figure 4.a) and arcs $\{v_3v_4,v_4v_3,v_3(v_2\circ v_5),(v_1\circ v_6)v_4\}$. On the other hand, if we change the order of the elements in U, then $D(U;V)$ is given by the graph in figure 4.b.*



(a) (b)

FIGURE 4. Digraphs with four vertices.

It is not difficult to see that

$$L(D,X)(U;V)=\begin{bmatrix}0&-1&x_3&-1\\0&0&-1&x_4\\0&0&0&-1\\-1&0&0&0\end{bmatrix}\approx\begin{bmatrix}x_3&-1&-1&0\\-1&x_4&0&0\\0&-1&0&0\\0&0&0&-1\end{bmatrix}=L(D(\{v_1,v_2\};\{v_6,v_5\}),X)_{\{x_{v_2\circ v_5}=0,\,x_{v_1\circ v_6}=-1\}}.$$

$$\approx\begin{bmatrix}x_3&-1&-1&0\\-1&x_4&0&0\\0&0&0&-1\\0&-1&0&0\end{bmatrix}=L(D(\{v_2,v_1\};\{v_6,v_5\}),X)_{\{x_{v_2\circ v_6}=0,\,x_{v_1\circ v_5}=0\}}.$$

In a similar way, we can define the digraph $D[U,V]$ that satisfies that

$$L(D,X)[U;V]\approx L(D[U;V],X)_{\{x_{u_i\circ v_i}=-m_{(v_i,u_i)}\,|\,i=1,\dots,s\}}.$$

The next results calculate the invariant γ for the path using the previous results on the minors of a generalized Laplacian matrix.

Corollary 4.10. *Let P_n be the path with n vertices and $\mathbb{Z}\subseteq\mathcal{P}$, then $\gamma_{\mathcal{P}}(P_n)=n-1$.*

Proof. Firstly, by example 4.6 we have that $P_n(v_1; v_n) = C_{n-1} \setminus \{\overrightarrow{v_1v_2}, \overrightarrow{v_{n-1}v_1}\}$. Hence, using theorem 4.1 and lemma 4.7 $\det(L(P_n, X)(v_1; v_n)) = 1$ and therefore $\gamma_{\mathcal{P}}(P_n) = n - 1$. ☐

A crucial open question discussed in [22, section 4] and [25, section 4] states that the critical group is cyclic for almost all simple graphs.

Conjecture 4.11. [25, Conjecture 4.2] *Given $n \geq 1$ and $0 \leq p \leq 1$, let $G(n,p)$ be the simple random graph with n vertices and $\Pr[uv \in G] = p$. If $c > 1$ and $c \cdot \log(n)/n < p < 1 - o(\log(n)/n)$, then*

$$\lim_{n\to\infty} \Pr[K(G(n,p)) \cong \mathbb{Z}_k] = 1.$$

However, we conjecture that critical ideals behave in an opposite way than critical groups.

Conjecture 4.12. *Let G be a simple graph with n vertices and $\mathbb{Z} \subseteq \mathcal{P}$, then $\gamma_{\mathcal{P}}(G) = n - 1$ if and only if $G = P_n$.*

4.1. **The critical ideals of the cycle.**

In this subsection we will calculate the critical ideals of a cycle. Let C_n be the cycle with n vertices and let $V(C) = \{v_1, v_2, \ldots, v_n\}$ be its vertex set. To simplify the notation, we consider the vertices of C_n as the classes modulo n. That is, for instance, the $n+1$-th vertex of C_n is the vertex v_1. In corollary 4.5 the determinant of the generalized Laplacian of C_n was calculated. Therefore, the critical ideals of C_n whenever $1 \leq i \leq n-1$ are the only ones that remain to calculate. In order to simplify the notation we will write $\det(G, X)$ instead of $\det(L(G, X))$.

First we will prove that almost all the critical ideals of C_n are trivial, except for i is equal to $n-1$ and n. Clearly $I_n(C_n, X)$ is generated by the determinant of $L(C_n, X)$. We will give a minimal set of generators for $I_{n-1}(C_n, X)$ and after that we will give a reduced Gröbner basis for it.

Theorem 4.13. *If C_n is the cycle with n vertices, then $I_i(C_n, X) = \langle 1 \rangle$ for all $1 \leq i \leq n-2$. Moreover,*

$$F_i = \{\det(C_n \setminus \{v_{i+1}, v_{i-1}\}, X), \det(C_n \setminus \{v_i, v_{i+2}\}, X), 1 + \det(C_n \setminus \{v_i, v_{i+1}\}, X)\}$$

for all $1 \leq i \leq n$ is a minimal set of generators for $I_{n-1}(C_n, X)$.

Proof. Firstly, we will prove that $I_i(C_n, X) = \langle 1 \rangle$ for all $1 \leq i \leq n-2$. Let D be the digraph obtained from the cycle with $n-2$ vertices, C_{n-2}, when we delete the arcs $\overrightarrow{v_1v_2}$ and $\overrightarrow{v_{n-2}v_1}$. It is not difficult to see that

$$L(C_n, X)(\{1, n\}; \{n-1, n\}) \approx L(D, X)_{\{x_1=0\}}.$$

Applying theorem 4.1 to the digraph D we have that $|\det(L(C_n, X)(\{1, n\}; \{n-1, n\}))| = 1$ and therefore $I_i(C_n, X) = \langle 1 \rangle$ for all $1 \leq i \leq n-2$.

Now, we will turn our attention to the $n-1$-th critical ideal of C_n. For all $1 \leq i, j \leq n$, let $Q_{i,j} = L(C_n, X)(i; j)$ and

$$q_{i,j} = \begin{cases} -\det(Q_{i,j}) & \text{if } \operatorname{lc}(\det(Q_{i,j})) = -1, \\ \det(Q_{i,j}) & \text{otherwise.} \end{cases}$$

Since $L(C_n, X)$ is symmetric, $Q_{i,j} \approx Q_{j,i}$ and $q_{i,j} = q_{j,i}$. Clearly, the ideal $I_{n-1}(C_n, X)$ is generated by the n^2 minors of size $n-1$ of $L(C_n, X)$.

On the other hand, let $1 \leq i \leq n$, $0 \leq l \leq n-1$, P_i^l be the induced path of C_n of length l that begins in the vertex i and ends in the vertex $i + l - 1$ and $p_{i,l} = \det(P_i^l, X)$. Note that, if $l = 0$, then P_i^l is the empty graph. For technical convenience we will adopt the convention that $p_{i,0} = 1$ for all $1 \leq i \leq n$ and $p_{i,-s} = -p_{i-s+1,s-2}$ for all $1 \leq s \leq i$. Note that, taking $s = 1$ we get that $p_{i,-1} = -p_{i-1+1,1-2} = -p_{i,-1}$ and therefore $p_{i,-1} = 0$ for all $1 \leq i \leq n$.

We present a relation between the $q_{i,j}$'s and the $p_{i,l}$'s. This relation will be useful to prove the next claim.

Claim 4.14. *If* $1 \leq i \leq j \leq n$, *then*

$$q_{i,j} = p_{i+1,j-i-1} + p_{j+1,n-j+i-1}.$$

Proof. At first, if $j = i$, then clearly $Q_{i,i} = L(C_n, X)(i; i) = L(C_n \setminus v_i, X \setminus x_i) = L(P_{i+1}^{n-1}, X)$. Now, if $j = i + 1$, then by lemma 4.7

$$Q_{i,i+1} = L(C_n, X)(i; i+1) \approx L(D, X)_{\{x_{v_i \circ v_{i+1}} = -1\}},$$

where D is the digraph obtained from C_n by deleting the arcs $\overrightarrow{v_i v_{i-1}}, \overrightarrow{v_{i+2} v_{i+1}}$ and identifying the vertices i and $i + 1$. Since D has only one spanning directed 1-factor (a directed cycle) and all the other directed 1-factors of D are 1-factors of $D \setminus v_i \circ v_{i+1} \cong P_{n-2}$, applying theorem 4.1 to D we get that $\det(L(D, X))_{\{x_{v_i \circ v_{i+1}} = -1\}} = -p_{i+2,n-2} - 1$ and therefore $q_{i,i+1} = p_{i+1,0} + p_{i+2,n-2}$. Finally, if $j \neq i, i + 1$, then by lemma 4.7

$$L(C_n, X)(i; j) \approx L(D, X)_{\{x_{v_i \circ v_j} = 0\}},$$

where D is the digraph obtained from C_n by deleting the arcs $\overrightarrow{v_i v_{i+1}}, \overrightarrow{v_{j-1} v_j}, \overrightarrow{v_i v_{i-1}}, \overrightarrow{v_{j+1} v_j}$ and identifying the vertices i and j. Since D has only two directed cycles containing the vertex $v_i \circ v_j$, applying theorem 4.1 we get the result. $\square$

Now, we present an algebraic relation between the $n - 1$-minors of $L(C_n, X)$ that will helpful to find a minimal set of generators for $I_{n-1}(C_n, X)$.

Claim 4.15. *If* $1 \leq i < j \leq n$ *and* $i \leq j + s \leq n + i$, *then*

$$q_{i,j+s} = p_{j+1,s-1} \cdot q_{i,j+1} - p_{j+2,s-2} \cdot q_{i,j}.$$

Proof. Firstly, if s is equal to 0 or 1, then the result is trivial. For the rest we will use induction on s. The proof will be divided in two cases, when $s \geq 2$ and $s \leq -1$. First assume that $s \geq 2$. Expanding the determinant of $L(P_s, X)$ we have that $p_{i+1,j-i} = x_j \cdot p_{i+1,j-i-1} - p_{i+1,j-i-2}$ and $p_{j,n-j+i} = x_j \cdot p_{j+1,n-j+i-1} - p_{j+2,n-j+i-2}$. Therefore

$$\begin{aligned} x_j q_{i,j} &\overset{4.14}{=} x_j \cdot [p_{i+1,j-i-1} + p_{j+1,n-j+i-1}] = p_{i+1,j-i} + p_{i+1,j-i-2} + p_{j,n-j+i} + p_{j+2,n-j+i-2} \\ &= p_{i+1,j-i-2} + p_{j,n-j+i} + p_{i+1,j-i} + p_{j+2,n-j+i-2} = q_{i,j-1} + q_{i,j+1}, \end{aligned}$$

and we get the result for $s = 2$. Now, if $s \geq 3$, then using induction hypothesis we get

$$\begin{aligned} q_{i,j+s+1} &= x_{j+s} \cdot q_{i,j+s} - q_{i,j+s-1} \\ &= x_{j+s} \cdot (p_{j+1,s-1} \cdot q_{i,j+1} - p_{j+2,s-2} \cdot q_{i,j}) - (p_{j+1,s-2} \cdot q_{i,j+1} - p_{j+2,s-3} \cdot q_{i,j}) \\ &= (x_{j+s} \cdot p_{j+1,s-1} - p_{j+1,s-2}) \cdot q_{i,j+1} - (x_{j+s} \cdot p_{j+2,s-2} - p_{j+2,s-3}) \cdot q_{i,j} \\ &= p_{j+1,s} \cdot q_{i,j+1} - p_{j+2,s-1} \cdot q_{i,j} \end{aligned}$$

We can use similar arguments when $s \leq -1$. $\square$

Claim 4.15 says us that if we fix i and j, then for all $1 \leq t \leq n$, $q_{i,t}$ can be generated by $q_{i,j}$ and $q_{i,j+1}$. Thus $\langle \{q_{i,t}\}_{t=1}^n \rangle = \langle q_{i,j}, q_{i,j+1} \rangle$ for all $1 \leq i, j \leq n$. Therefore it is not difficult to see that

$$I_{n-1}(C_n, X) = \langle q_{i,i+1}, q_{i,i+2}, q_{i-1,i+1} \rangle \text{ for all } 1 \leq i \leq n.$$

$\square$

Before to present a Gröbner basis for $I_{n-1}(C_n, X)$ we present a identity between the $q_{i,j}$'s. This identity will be very useful to reduce the S-polynomials.

Proposition 4.16. *If* $1 \leq i \leq n$ *and* $0 \leq k, s \leq n-i$, *then*

$$p_{i+k+1,s} \cdot q_{i,i+k+1} - p_{i+1,s} \cdot q_{i+s,i+k+s+1} = p_{i+k+2,s-1} \cdot q_{i,i+k} - p_{i+1,s-1} \cdot q_{i+s+1,i+k+s+1}.$$

Proof. Applying claim 4.15 with $i' = i$, $j' = i+k$, and $s' = s+1$ we get that

$$q_{i,i+k+s+1} = q_{i',j'+s'} = p_{j'+1,s'-1} \cdot q_{i',j'+1} - p_{j'+2,s'-2} \cdot q_{i',j'} = p_{i+k+1,s} \cdot q_{i,i+k+1} - p_{i+k+2,s-1} \cdot q_{i,i+k}.$$

Also, applying claim 4.15 with $i' = i+k+s+1$, $j' = i+s$, $s' = -s$, and using that $q_{i,j} = q_{j,i}$ and $p_{i,-s} = -p_{i-s+1,s-2}$ we get that

$$\begin{aligned} q_{i,i+k+s+1} &= q_{i+k+s+1,i} = q_{i',j'+s'} = p_{j'+1,s'-1} \cdot q_{i',j'+1} - p_{j'+2,s'-2} \cdot q_{i',j'} \\ &= p_{i+s+1,-(s+1)} \cdot q_{i+k+s+1,i+s+1} - p_{i+s+2,-(s+2)} \cdot q_{i+k+s+1,i+s} \\ &= p_{i+1,s} \cdot q_{i+s,i+k+s+1} - p_{i+1,s-1} \cdot q_{i+s+1,i+k+s+1}. \end{aligned}$$

Therefore

$$p_{i+k+1,s} \cdot q_{i,i+k+1} - p_{i+k+2,s-1} \cdot q_{i,i+k} = q_{i,i+k+s+1} = p_{i+1,s} \cdot q_{i+s,i+k+s+1} - p_{i+1,s-1} \cdot q_{i+s+1,i+k+s+1}$$

and we get the result. ☐

Now, we are ready to find a Gröbner basis for $I_{n-1}(C_n, X)$. We divide this in odd and even cases.

Theorem 4.17. *Let* $n = 2m+1$ *and*

$$B_1 = \{q_{i,i+m+1} \,|\, i = 1, \dots, n\} = \{q_{i,i+m} \,|\, i = 1, \dots, n\}.$$

Then B_1 *is a reduced Gröbner basis for* $I_{n-1}(C_n, X)$ *with respect to the graded lexicographic order with* $x_1 > \cdots > x_n$.

Proof. Note that, $\mathrm{lt}(q_{i,i+m+1}) \overset{4.14}{=} \mathrm{lt}(p_{i+1,m}) = x_{i+1} \cdots x_{i+m}$ and therefore $\mathrm{lt}(q_{i,i+m+1}) \nmid \mathrm{lt}(q_{i',i'+m+1})$ for all $1 \leq i < i' \leq n$. That is, if B_1 is a Gröbner basis for $I_{n-1}(C_n, X)$, then it is reduced.

Now, in order to prove that B_1 is a Gröbner basis for $I_{n-1}(C_n, X)$ we need to show that all the S-polynomials of the elements on B_1 can be reduced to 0 by elements on B_1. Since

$$S(q_{i,i+m+1}, q_{i',i'+m+1}) = -S(q_{i',i'+m+1}, q_{i,i+m+1}),$$

we can assume without loss of generality that $i < i'$. It is not difficult to see that

$$\begin{aligned} S(q_{i,i+m+1}, q_{i+s,i+m+s+1}) &= (x_{i+m+1} \cdots x_{i+m+s}) \cdot q_{i,i+m+1} - (x_{i+1} \cdots x_{i+s}) \cdot q_{i+s,i+m+s+1} \\ &= \mathrm{lt}(p_{i+m+1,s}) \cdot q_{i,i+m+1} - \mathrm{lt}(p_{i+1,s}) \cdot q_{i+s,i+m+s+1} \end{aligned}$$

for all $1 \leq i \leq n-1$ and $1 \leq s \leq n-i$. By proposition 4.16 with $k = m$ we get that

$$\begin{aligned} S(q_{i,i+m+1}, q_{i+s,i+m+s+1}) &= -(p_{i+m+1,s} - \mathrm{lt}(p_{i+m+1,s})) \cdot q_{i,i+m+1} + (p_{i+1,s} - \mathrm{lt}(p_{i+1,s})) \cdot q_{i+s,i+m+s+1} \\ &\quad + p_{i+m+2,s-1} \cdot q_{i,i+m} - p_{i+1,s-1} \cdot q_{i+s+1,i+m+s+1}. \end{aligned}$$

Finally, since $\mathrm{lt}(q_{i,i+m+1}) = \mathrm{lt}(p_{i+1,m})$, $\mathrm{lt}(q_{i,i+m}) = \mathrm{lt}(p_{i+m+1,m})$, $\mathrm{lt}(p_{i+1,s-1}) = x_{i+1} \cdots x_{i+s-1}$, and

$$\mathrm{lt}(p_{i+1,s} - \mathrm{lt}(p_{i+1,s})) = \begin{cases} x_{i+1} \cdots x_{i+s-2} & \text{if } 1 \notin \{i+1, \cdots, i+s\}, \\ x_1 \cdots x_{i+s} x_{i+1} \cdots x_{n-2} & \text{otherwise,} \end{cases}$$

for all $1 \leq i, s \leq n$. Then $S(q_{i,i+m+1}, q_{i+s,i+s+m+1}) \rightarrow_{B_1} 0$ for all $1 \leq i \leq n-1$ and $1 \leq s \leq n-i$. ☐

The even case is slightly different.

Theorem 4.18. *Let $n = 2m$ and*

$$B_0 = \{q_{i,i+m} \,|\, i = 0, \dots, m-1\} \cup \{q_{i+m,i+1} \,|\, i = 0, \dots, m-2\}.$$

Then B_0 is a Gröbner basis for $I_{n-1}(C_n, X)$ with respect to the graded lexicographic order with $x_{m-1} > \cdots > x_{2m} > x_1 \cdots > x_{m-2}$.

Proof. Following the proof of theorem 4.17, we have that $\mathrm{lt}(q_{i,i+m}) = \mathrm{lt}(p_{i+1,m-1}) = x_{i+1} \cdots x_{i+m-1}$ for all $i = 0, \dots, m-1$ and $\mathrm{lt}(q_{i+m,i+1}) = \mathrm{lt}(p_{i+m+1,m}) = x_{i+m+1} \cdots x_i$ for all $i = 0, \dots, m-2$. Therefore $\mathrm{lt}(q_{i,i+m}) \not| \mathrm{lt}(q_{i',i'+m})$ for all $1 \leq i < i' \leq m-1$, $\mathrm{lt}(q_{i+m,i+1}) \not| \mathrm{lt}(q_{i'+m,i'+1})$ for all $1 \leq i < i' \leq m-2$, and $\mathrm{lt}(q_{i,i+m}) \not| \mathrm{lt}(q_{i'+m,i'+1})$ for all $1 \leq i \leq m-1$ and $1 \leq i' \leq m-2$. Moreover, since $\deg(q_{i+m,i+1} - \mathrm{lt}(q_{i+m,i+1})) = m-2$ and $\mathrm{lt}(q_{i,i+m} - \mathrm{lt}(q_{i,i+m})) = \mathrm{lt}(p_{i+m+1,m-1}) = x_{i+m+1} \cdots x_{i-1}$, if B_0 is a Gröbner basis for $I_{n-1}(C_n, X)$, then it is reduced.

In a similar way to the proof of theorem 4.17, we need to show that all the S-polynomials of the elements on B_0 can be reduced to 0 by elements on B_0. At difference to the proof of theorem 4.17, in this case we have three types of S-polynomials of elements of B_0. First

$$\begin{aligned} S(q_{i,i+m}, q_{i+s,i+m+s}) &= (x_{i+m} \cdots x_{i+m+s-1}) \cdot q_{i,i+m} - (x_{i+1} \cdots x_{i+s}) \cdot q_{i+s,i+m+s} \\ &= \mathrm{lt}(p_{i+m,s}) \cdot q_{i,i+m} - \mathrm{lt}(p_{i+1,s}) \cdot q_{i+s,i+m+s} \end{aligned}$$

for all $0 \leq i \leq m-2$ and $1 \leq s \leq m-i-1$. Applying proposition 4.16 with $k = m-1$ we get that

$$\begin{aligned} S(q_{i,i+m}, q_{i+s,i+m+s}) &= -(p_{i+m,s} - \mathrm{lt}(p_{i+m,s})) \cdot q_{i,i+m} + (p_{i+1,s} - \mathrm{lt}(p_{i+1,s})) \cdot q_{i+s,i+m+s} \\ &\quad + p_{i+m+1,s-1} \cdot q_{i,i+m-1} - p_{i+1,s-1} \cdot q_{i+s+1,i+m+s}. \end{aligned} \tag{4.1}$$

When $i \geq 1$ and $i + s \leq m-2$ is not difficult to use this identity to prove that $S(q_{i,i+m}, q_{i+s,i+m+s})$ it reduces to 0 by B_0. Now, only remain to analyze two special cases, when $i = 0$ or $i + s = m-1$. Taking $i = 0$ in equation 4.1 and the fact that

$$q_{0,m-1} = q_{m-1,2m} \overset{4.15}{=} p_{2m-1,1} \cdot q_{m-1,2m-1} - p_{2m,0} q_{m-1,2m-2} = x_{2m-1} \cdot q_{m-1,2m-1} - q_{m-1,2m-2}$$

we have that

$$\begin{aligned} S(q_{0,m}, q_{s,m+s}) &= -(p_{m,s} - \mathrm{lt}(p_{m,s})) \cdot q_{0,m} + (p_{1,s} - \mathrm{lt}(p_{1,s})) \cdot q_{s,m+s} + x_{2m-1} \cdot p_{m+1,s-1} \cdot q_{m-1,2m-1} \\ &\quad - \begin{cases} p_{m+1,s-1} \cdot q_{m-1,2m-2} + p_{1,s-1} \cdot q_{s+1,m+s} & \text{if } s \leq m-3, \\ (p_{m+1,m-3} + p_{1,m-3}) \cdot q_{m-1,2m-2} & \text{if } s = m-2. \end{cases} \end{aligned}$$

Therefore $S(q_{0,m}, q_{s,m+s}) \to_{B_0} 0$ for all $1 \leq s \leq m-2$. On the other hand, taking $i + s = m-1$ in equation 4.1 and using that $q_{m,2m-1} = q_{2m-1,m} \overset{4.15}{=} x_{2m} \cdot q_{0,m} - q_{1,m}$ we have that

$$\begin{aligned} S(q_{i,i+m}, q_{m-1,2m-1}) &= -(p_{i+m,s} - \mathrm{lt}(p_{i+m,s})) \cdot q_{i,i+m} + (p_{i+1,s} - \mathrm{lt}(p_{i+1,s})) \cdot q_{m-1,2m-1} \\ &\quad + p_{i+m+1,s-1} \cdot q_{i,i+m-1} - p_{i+1,s-1} \cdot q_{m,2m-1} \\ &= -(p_{i+m,s} - \mathrm{lt}(p_{i+m,s})) \cdot q_{i,i+m} + (p_{i+1,s} - \mathrm{lt}(p_{i+1,s})) \cdot q_{m-1,2m-1} \\ &\quad + \begin{cases} (p_{m+2,s-1} + p_{2,s-1}) \cdot q_{1,m} - x_{2m} \cdot p_{2,s-1} \cdot q_{0,m} & \text{if } i = 1, \\ p_{i+m+1,s-1} \cdot q_{i,i+m-1} - x_{2m} \cdot p_{i+1,s-1} \cdot q_{0,m} + p_{i+1,s-1} \cdot q_{1,m} & \text{if } i \geq 2. \end{cases} \end{aligned}$$

Therefore, $S(q_{i,i+m}, q_{m-1,2m-1})$ can be reduced to 0 by B_0. Finally, if $i = 0$ and $s = m-1$ we have that

$$S(q_{0,m}, q_{m-1,2m-1}) = (q_{0,m} - \mathrm{lt}(q_{0,m}))q_{m-1,2m-1} - (q_{m-1,2m-1} - \mathrm{lt}(q_{m-1,2m-1}))q_{0,m}.$$

Now, we follow with the S-polynomials of a second type.

$$\begin{aligned} S(q_{i+m,i+1}, q_{i+m+s,i+s+1}) &= (x_{i+1}\cdots x_{i+s})\cdot q_{i+m,i+1} - (x_{i+m+1}\cdots x_{i+m+s})\cdot q_{i+m+s,i+s+1} \\ &= \mathrm{lt}(p_{i+1,s})\cdot q_{i+m,i+1} - \mathrm{lt}(p_{i+m+1,s})\cdot q_{i+m+s,i+s+1} \end{aligned}$$

for all $0 \leq i \leq m-3$ and $1 \leq s \leq m-i-2$. By proposition 4.16 with $i = i+m$ and $k = -m$ we get that

$$\begin{aligned} S(q_{i+m,i+1}, q_{i+m+s,i+s+1}) &= -(p_{i+1,s} - \mathrm{lt}(p_{i+1,s}))\cdot q_{i+m,i+1} + (p_{i+m+1,s} - \mathrm{lt}(p_{i+m+1,s}))\cdot q_{i+m+s,i+s+1} \\ &\quad + p_{i+2,s-1}\cdot q_{i+m,i} - p_{i+m+1,s-1}\cdot q_{i+m+s+1,i+s+1} \end{aligned}$$

for all $0 \leq i \leq m-3$ and $1 \leq s \leq m-i-2$. Therefore, $S(q_{i+m,i+1}, q_{i+m+s,i+s+1}) \rightarrow_{B_0} 0$. We finish with the S-polynomials of a third type.

$$\begin{aligned} S(q_{i,i+m}, q_{i'+m,i'+1}) &= \begin{cases} (x_{i'+m+1}\cdots x_i)\cdot q_{i,i+m} - (x_{i'+1}\cdots x_{i+m-1})\cdot q_{i'+m,i'+1} & \text{if } i \leq i'-1, \\ (x_{i'+m+1}\cdots x_{i'})\cdot q_{i,i+m} - (x_{i+1}\cdots x_{i+m-1})\cdot q_{i'+m,i'+1} & \text{if } i = i', \\ (x_{i'+m+1}\cdots x_{i'})\cdot q_{i,i+m} - (x_{i+1}\cdots x_{i+m-1})\cdot q_{i'+m,i'+1} & \text{if } i = i'+1, \\ (x_{i+m}\cdots x_{i'})\cdot q_{i,i+m} - (x_{i+1}\cdots x_{i'+m})\cdot q_{i'+m,i'+1} & \text{if } i \geq i'+2, \end{cases} \\ &= \begin{cases} \mathrm{lt}(p_{i'+m+1,m+i-i'})\cdot q_{i,i+m} - \mathrm{lt}(p_{i'+1,m+i-i'-1})\cdot q_{i'+m,i'+1} & \text{if } i \leq i', \\ \mathrm{lt}(p_{i+m,m+i'-i+1})\cdot q_{i,i+m} - \mathrm{lt}(p_{i+1,m+i'-i})\cdot q_{i'+m,i'+1} & \text{if } i \geq i'+1, \end{cases} \end{aligned}$$

for all $0 \leq i \leq m-1$ and $0 \leq i' \leq m-2$. If $i = i', i'+1$, then

$$S(q_{i,i+m}, q_{i'+m,i'+1}) = (q_{i,i+m} - \mathrm{lt}(q_{i,i+m}))q_{i'+m,i'+1} - (q_{i'+m,i'+1} - \mathrm{lt}(q_{i'+m,i'+1}))q_{i,i+m}.$$

Using similar arguments to those in proposition 4.16 is not difficult to prove that

$$p_{i'+m+1,m+i-i'}\cdot q_{i,i+m} - p_{i'+1,m+i-i'-1}\cdot q_{i'+m,i'+1} = p_{i'+m+1,m+i-i'-1}\cdot q_{i+1,i+m} - p_{i'+2,m+i-i'-2}\cdot q_{i'+m,i'},$$

$$p_{i+m,m+i'-i+1}\cdot q_{i,i+m} - p_{i+1,m+i'-i}\cdot q_{i'+m,i'+1} = p_{i+m+1,m+i'-i}\cdot q_{i,i+m-1} - p_{i+1,m+i'-i-1}\cdot q_{i'+m+1,i'+1}.$$

Finally, using this identities it can be proved that $S(q_{i,i+m}, q_{i'+m,i'+1}) \rightarrow_{B_0} 0$ for all $0 \leq i \leq m-1$ and $0 \leq i' \leq m-2$. □

Example 4.19. *Let* $n = 2m = 2(3) = 6$*, then* $q_{6,3} = \underline{x_1x_2} + x_4x_5 - 2$, $q_{1,4} = \underline{x_2x_3} + x_5x_6 - 2$, $q_{2,5} = \underline{x_3x_4} + x_1x_6 - 2$, $q_{3,1} = \underline{x_4x_5x_6} - x_4 - x_6 + x_2$, $q_{4,2} = \underline{x_1x_5x_6} - x_1 - x_5 + x_3$ *(the leading terms are underlined) forms a Gröbner basis for* $I_5(C_6, X)$ *with respect to the graded lexicographic order with* $x_2 > x_3 > x_4 > x_5 > x_6 > x_1$*. For instance, taking* $i = 0$ *and* $s = m-1 = 2$ *it is not difficult to see that*

$$S(q_{6,3}, q_{2,5}) = x_3x_4q_{6,3} - x_1x_2q_{2,5} = (x_4x_5 - 2)q_{2,5} - (x_5x_6 - 2)q_{3,6}.$$

Also, taking $i = 1$ *and* $s = 1$ *we get that*

$$S(q_{1,4}, q_{2,5}) = x_4q_{1,4} - x_2q_{2,5} = x_4x_5x_6 - 2x_4 - x_2x_1x_6 + 2x_2 = q_{3,1} - q_{3,5} = 2q_{3,1} - x_6q_{6,3}.$$

In order to finish this article we will present a simple application of theorem 4.13, see remark 6.8 in [22] for a similar result for graphs.

Corollary 4.20. *Let* D *be a digraph and* $v \in V(D)$ *such that* $D \setminus v \simeq C_n$*, then* $K(D)$ *has at most two invariant factors different to one.*

It is not easy to determine when the critical group of G is cyclic because in order to do this, we need to find the greatest common divisor of the evaluation of polynomials.

Acknowledgments

The authors would like to thank the anonymous referees for their helpful comments.

References

[1] W. W. Adams and P. Loustaunau, An introduction to Gröbner basis, GSM-AMS (1994).

[2] C. A. Alfaro and C. E. Valencia, Graphs with a large number of invariant factors, manuscript.

[3] C. A. Alfaro and C. E. Valencia, Graphs with $\gamma = 3$, in preparation.

[4] H. Bai, On the critical group of the n-cube, Linear Algebra Appl. 369 (2003) 251–261.

[5] J. Bang-Jensen and G. Gutin, Digraphs: Theory, Algorithms and Applications, Springer-Verlag, London, 2006.

[6] P. Bak, C. Tang, and K. Wiesenfeld, Self-organized criticality: an explanation of the 1/f noise. Phys. Rev. Lett. 59 (1987), no. 4, 381–384.

[7] A. Berget, A. Manion, M. Maxwell, A. Potechin, and V. Reiner, The critical group of a line graph, Annals of Combinatorics 16 / 3 (2012), 449–488.

[8] N. L. Biggs, Algebraic Graph Theory, second ed., Cambridge Mathematical Library, Cambridge University Press, Cambridge, 1993.

[9] H. Christianson and V. Reiner, The critical group of a threshold graph, Linear Algebra Appl. 349 (2002) 233–244.

[10] R. Cori and D. Rossin, On the sandpile group of dual graphs, Europ. J. Combinatorics 21 (2000) 447–459.

[11] R. Cori, D. Rossin, and B. Salvy, Polynomial ideals for sandpiles and their Göbner bases, Theoretical Computer Science, 276 (2002) 1–15.

[12] H. Corrales and C. Valencia, On the critical ideals of trees, in preparation.

[13] H. Corrales and C. Valencia, On the critical group of matrices, Bulletin Mathématique de la Socité des Sciences Mathématiques de Roumanie **54** (102) (2011) 213–236

[14] D. Dhar, Self-organized critical state of sandpile automaton models. Phys. Rev. Lett. 64 (1990), no. 14 1613–1616.

[15] R. Diestel, Graph Theory, Graduate Texts in Mathematics 173, Springer-Verlag, New York, 2005.

[16] A. M. Duval, C. J. Klivans, and J. L. Martin, Critical groups of simplicial complex, to appear in Annals of Combinatorics, `arXiv:1101.3981v2 [math.CO]`.

[17] C. D. Godsil and G. F. Royle, Algebraic Graph Theory, Graduate Texts in Mathematics 207, Springer-Verlag, New York, 2001.

[18] B. Jacobson, A. Niedermaier, and V. Reiner, Critical groups for complete multipartite graphs and Cartesian products of complete graphs, J. Graph Theory 44 (2003) 231–250.

[19] L. Levine, The sandpile group of a tree, Europ. J. Combinatorics 30 (2009), no. 4 1026–1035.

[20] D. J. Lorenzini, Arithmetical graphs, Math. Ann. 285 (1989) 481–501.

[21] D. J. Lorenzini, A finite group attached to the Laplacian of a graph, Discrete Math. 91 (1991) 277–282.

[22] D. J. Lorenzini, Smith normal form and Laplacians, J. Combin. Theory B 98 (2008), 1271–1300.

[23] E. R. Speer, Asymmetric Abelian sandpiles models, J. Statistical Phys. 71 1-2 (1993) 61-74.

[24] E. Toumpakari, On the sandpile group of regular trees, Europ. J. Combin. 28 (2007) 822–842.

[25] D. G. Wagner, The critical group of a directed graph, preprint, 2000. `ArXiv:math/0010241`.

Departamento de Matemáticas, Centro de Investigación y de Estudios Avanzados del IPN, Apartado Postal 14–740, 07000 Mexico City, D.F.

E-mail address, Hugo Corrales: `hhcorrales@gmail.com`

Departamento de Matemáticas, Centro de Investigación y de Estudios Avanzados del IPN, Apartado Postal 14–740, 07000 Mexico City, D.F.

E-mail address, Carlos E. Valencia[1]: `cvalencia@math.cinvestav.edu.mx, cvalencia75@gmail.com`

---

[1] Corresponding author